

**SECTION 18 EXEMPTION USE DIRECTIONS
FOR USE AND DISTRIBUTION ONLY WITHIN ADA, CANYON, GEM, NAMPA, OWYHEE, PAYETTE, TWIN
FALLS AND WASHINGTON COUNTIES OF IDAHO
TO CONTROL MITES IN SWEET CORN GROWN FOR SEED**

Onager
Miticide



EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY

**FILE SYMBOL NUMBER: 09-ID-04
EPA REG. NO: 10163-277
EFFECTIVE DATE: June 29, 2009
EXPIRATION DATE: August 20, 2009**

ACTIVE INGREDIENT: Hexythiazox	% By Wt.
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide	11.8%
INERT INGREDIENTS	88.2%
	TOTAL 100.0%

THIS LABEL IS IN ADDITION TO THE STANDARD LABEL PRINTED ON THE CONTAINER

GENERAL INFORMATION

Onager controls mites through its activity on eggs and immature stages. Onager does not kill adult mites although treated females will fail to lay viable eggs while in contact with treated foliage. Onager is not systemic and thorough coverage of leaf surfaces is necessary for effective control.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable restrictions, precautions, worker protection standard requirements, and directions for use on the EPA Section 3 label must be followed.
- This labeling should be in the possession of the user at the time of pesticide application.

Application Equipment

Ground Application: Apply the recommended rate of Onager by ground equipment at 15-20 GPA. Use a spray boom equipped to position one 16-inch boom drop extension per row, with two nozzles directed to spray up and under the crop canopy – one to the right and one to the left.

Air Application: Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.

Chemigation: Apply Onager only through sprinkler, including center pivot, lateral move, Low Energy Precision Applications (LEPA), end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply Onager through any other type of irrigation system.

CROP	PEST	RATE Oz./acre	COMMENTS
Sweet Corn Grown for Seed	Banks grass mite; Two spotted spider mite	10-24	Apply Onager at the first sign of mites before populations begin to build.
	<ul style="list-style-type: none"> • Do not apply more than a total of 24 oz. formulated product per acre per year. • Do not make more than one application per year. • Do not apply within 45 days of harvest. 		

Applications of Onager under this Section 18 Emergency Specific Exemption are limited to infestations that meet the following Economic Threshold Criteria: Presence of mites on sweet corn foliage or on wheat or grasses in or adjacent to the cornfield.

Environmental Hazards: This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.