

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®] Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN ID-090003
For Distribution and Use Only in the State of Idaho

Control of Billbugs, Cutworms and Aphids Infesting Perennial Grass Seed Crops During the Year of Establishment

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Billbugs and Cutworms: Apply 1 quart of Lorsban Advanced per acre in 20 gallons of water per acre as a broadcast spray using power-operated ground spray equipment. For aerial application, apply in 2 to 5 gallons of water per acre. For improved results, thatch should be moist at time of application and the insecticide should be washed into the thatch immediately after application. If sprinkler irrigation is available, apply 1/4 to 1/2 inch of water after application to help incorporate the material, or apply to early morning dew-covered fields, or during or immediately following a light rain. For **billbug control**, apply in early April when adult billbugs are active. For **cutworm control**, apply in either spring or fall when larvae or larval damage first appears. Fall applications should follow sufficient irrigation and/or rainfall.

Aphids: Apply 1 to 2 pints of Lorsban Advanced per acre in 20 gallons of water per acre as a broadcast spray using power-operated ground spray equipment. For aerial application, use 2 to 5 gallons of water per acre. Treat before heading as colonies begin to form.

Consult your cooperative agricultural extension service for additional information concerning billbug, cutworm, and aphid control practices in your area.

Specific Use Precautions and Restrictions:

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than three applications of Lorsban Advanced per year.
- Maximum single application rate is 0.94 lb ai chlorpyrifos per acre (maximum of 2.82 lb ai chlorpyrifos per year).
- Use of Lorsban Advanced according to this supplemental label is deemed a non-food use by the Idaho Department of Agriculture and, as such, is not for use on fields producing grass for livestock feed.
- Do not feed or graze grass following application of Lorsban Advanced.
- Do not cut treated grass for forage, hay or silage.
- No portion of the treated field may be used for human or animal feed.
- Lorsban Advanced may only be applied to grass seed fields during the year of establishment.

- Harvest of grass seed from fields treated with Lorsban Advanced cannot occur for at least 365 days from the date of application.
- **Chemigation:** Do not apply through any type of irrigation equipment.
- Growers may not treat the same perennial grass seed crop during the year of establishment under the requirements of any SLN for Lorsban-4E.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Broadleaf weed bloom must be controlled prior to the application in grass seed fields with 5 blooms or more per square yard.
- This product is toxic to fish, aquatic invertebrates, small mammals and birds, and should not be used where impact on listed threatened or endangered species is likely. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. Contact the Idaho Department of Fish & Wildlife, National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species (e.g., bull trout, chinook salmon). Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

This label expires and must not be used or distributed after February 3, 2014.

®Trademark of Dow AgroSciences LLC

R368-036

Approved: 02/03/09

Initial printing.