

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban[®] Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN ID-090004

For Distribution and Use Only in the State of Idaho

Control of Grape Mealybugs and Cutworms Infesting Grapes

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Refer to product label for Use Precautions and Restrictions, Mixing and Application instructions.

Cutworms: Apply Lorsban Advanced at the rate of 1 quart per acre as a banded application to the base of the vines in a minimum spray volume of at least 50 gallons of water using power-operated ground spray equipment. Treat when cutworms first become active and when field counts indicate damaging insect populations are developing or present. Do not apply after bloom stage of growth. Consult your state agricultural experiment station or extension service specialist concerning cutworm control practices in your area.

Grape Mealybugs: Apply Lorsban Advanced at the rate of 1 to 2 quarts per acre in a minimum spray volume of at least 50 gallons of water per acre using ground equipment only. To obtain thorough coverage, spray all above-ground parts, especially the trunk and cordons where mealybugs are located. Use the 1 quart rate for normal infestations prior to bloom stage of growth. Use the 2 quart rate for high infestations as a dormant or late dormant application only. If an application is made at the 2 quart rate, a second application is not allowed.

Specific Use Precautions and Restrictions:

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- **Preharvest Interval:** Do not apply within 35 days of harvest.
 - Do not apply more than 2 quarts of Lorsban Advanced (1.88 lb ai chlorpyrifos) per acre per crop growing season or make more than two applications. If the 2 quart application rate is used, only one application is allowed.
 - **Chemigation:** Do not apply through any type of irrigation equipment.
 - Applications after budbreak may result in transient leaf yellowing (Concords).

This label expires and must not be used or distributed after February 3, 2014.

[®]Trademark of Dow AgroSciences LLC

R368-060

Approved: ___/___/___

Replaces R368-037.