

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



For Agricultural or Commercial Use Only
EMERGENCY CALLS: 800-331-3148

24(c) SPECIAL LOCAL NEED LABEL
SUGARBEET

FOR DISTRIBUTION AND USE ONLY IN
IDAHO

IT IS A VIOLATION OF FEDERAL LAW
TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH IT LABELING

EPA SLN No. ID-040004

This label for Mustang Max is valid until December 31, 2008 or until otherwise amended, withdrawn, canceled or suspended.

CROP	PEST	RATE OF APPLICATION
SUGAR BEET (At Plant)	Seedling Pests including: Wireworm (Suppression Only) White Grub (Suppression Only) Cutworms species	4.0 ounces per acre (0.025 lb ai)
SUGAR BEET (Foliar)	Foliar Pests including: Cutworms species Flea Beetle Sugar Beet Root Maggot Adults	2.24 to 4.0 ounces per acre (0.014 - 0.025 lb ai)

Directions For Use

At Plant Applications

For wireworm and white grub, apply in-furrow or in a 3 to 4 inch T-band over the open furrow at planting in a minimum of 3 to 5 gallons of water per acre at the rate of 4 ounces per acre.

For cutworm species, apply at planting on the soil surface in a 5 to 7 inch band or broadcast in a minimum of 3 to 5 gallons per acre at the rate of 4 ounces per acre.

Foliar Applications :

Apply Mustang Max at the rate of 2.24 to 4 ounces per acre. For control of sugar beet root maggot adults apply at the rate of 4 ounces per acre. A second application 1-2 weeks later may be needed.

Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and, or scouting results.

Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. Use a minimum of 3 gallons per acre by air and 10 gallons per acre by ground.

Applications may be made by chemigation, refer to the Mustang Max Section 3 label for instructions.

Restrictions

Observe all spray drift precautions on the EPA registered label.

Do not apply within 50 days of harvest of sugar beet tops or roots.

Do not apply more than 0.075 pound active ingredient per acre per season including at plant plus foliar applications of Mustang Max.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED
MUSTANG MAX LABEL (EPA Registration No 279-3249) MUST BE FOLLOWED.**

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



24(c) Registrant
FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

MustangMAX_24c_ID_sugarbeet_12-14-2006

Mustang Max and **FMC** -FMC Trademarks
©2006 FMC Corporation