

Supplemental Labeling

RECEIVED

SEP 02 2008

DIV. OF AG. RESOURCES



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

EPA 24(c) Special Local Need Registration SLN #ID-080015
For Distribution and Use Only Within the State of Idaho

Control of Powdery Mildew (*Erysiphe heraclei*) in Carrots Grown for Seed

Special Conditions and Risks of Use

USE OF Rally 40WSP (THE "PRODUCT") ON CARROTS GROWN FOR SEED (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Dow AgroSciences makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;
- (3) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Dow AgroSciences, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Dow AgroSciences and not specified by the U.S. EPA or the State of Idaho.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

Rally 40WSP is a systemic, protectant and curative fungicide. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program, starting at the first sign of disease development and continuing until the threat of further disease development is past.

Apply 5 to 7.5 oz of Rally 40WSP (0.125 to 0.187 lb active ingredient) in a minimum of 20 gallons of water per acre for ground application or in a minimum of 10 gallons of water per acre for aerial application. Begin applications prior to or at the start of disease development and continue applications on a 14- to 21-day application schedule. Use higher rate or shorter spray interval on susceptible varieties or under heavy disease pressure.

Specific Use Restrictions:

- Do not apply more than 15 oz of Rally 40WSP (0.375 lb active ingredient) per acre per year.
- **Chemigation:** Do not apply this product through any type of irrigation system for the use described in this supplemental labeling.
- **Rotation Crops:** Fields treated with Rally 40WSP can be rotated at any time to crops that are included on the product label. All other crops may be planted 30 days after the final application of Rally 40WSP.
- Use of Rally 40WSP according to this supplemental label is deemed a non-food use by the Idaho Department of Agriculture and, as such, is not for use on fields producing carrots for livestock feed.
- Do not feed or graze carrots following application of Rally 40WSP.
- Do not cut treated carrots for hay or forage.
- Do not use harvested seed for sprouting.
- No portion of the treated field, including seed, seed screenings, hay forage, or stubble may be used for human or animal feed.

®Trademark of Dow AgroSciences LLC

R182-029

Accepted: ___/___/___

Initial printing.