



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina
 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

**SENCOR® DF 75% DRY FLOWABLE
 HERBICIDE**

For use on potatoes.

EPA Reg. No. 264-738

EPA SLN. No. ID-870017

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Do not apply this product through any type of irrigation system.

APPLICATION INSTRUCTIONS		
CROP	PEST	DOSAGE SENCOR DF
Potatoes	Weeds	2/3 to 1-1/3 lbs/A
<p>REMARKS: PRE-PLANT INCORPORATED: Apply 2/3 to 1-1/3 lbs of SENCOR DF per acre as a broadcast spray or impregnated on dry bulk fertilizer. Incorporate 4-6" deep by double discing at right angles. Refer to the SENCOR label for proper rate for soil type.</p> <p>Dry bulk fertilizer may be impregnated or coated with SENCOR for pre-plant incorporated application to potatoes. Refer to the "Commercial Impregnation and Application of SENCOR on Dry Bulk Fertilizer" section of the label for information, instructions, and proper procedure. All recommendations, cautions, and special precautions on the SENCOR label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling. Apply pre-plant using 500-1200 lbs dry bulk fertilizer per acre.</p> <p>Before any fertilizer spreader is used, it must be calibrated. If a cyclone type spreader is used, the application must have two passes or a 100% overlap. Field ends and turn around areas may be over or under treated, resulting in a lack of weed control or possible crop injury.</p>		

NOTE: Use only on russeted or on white-skinned potatoes that are not early maturing. Do not use this method of treatment on red skinned potatoes or on Centennial, Atlantic, Shepody, Chipbelle, or Bellchip varieties or crop injury may result. Temporary chlorosis (yellowing) may occur under prolonged conditions of cool, wet or cloudy weather. All applicable directions, restrictions and precautions on the EPA-registered label for all products used are to be followed.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)
 or
 Visit our worldwide web site at www.bayercropscienceus.com

Sencor DF is a registered trademark of Bayer.