



**THIONEX® 3EC**  
**EPA REG. NO. 66222-63**  
**COOLEY SPRUCE ADELGID AND DOUGLAS**  
**FIR NEEDLE MIDGE**  
**IN CONIFERS (GROWN AS CHRISTMAS TREES)**  
**IN THE STATE OF IDAHO**  
**EPA SLN NO. ID-030002**

**SUPPLEMENTAL LABELING FOR DISTRIBUTION AND USE ONLY  
IN THE STATE OF IDAHO**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

Personal Protective Equipment and Reentry Intervals on the basic label must be followed.

CROP	PEST	RATE OF APPLICATION
Conifers (Grown as Christmas Trees)	Cooley Spruce Adelgid and Douglas Fir Needle Midge, Eriophyid Needle mite	$\frac{2}{3}$ quart per 100 gallons of water

**DIRECTIONS FOR USE**

Use  $\frac{2}{3}$  of a quart of Thionex 3EC (1/2 pound active ingredient) per 100 gallons of water. For air application, use  $\frac{2}{3}$  to  $1\frac{1}{3}$  quarts of Thionex 3EC (1/2 to 1 pound active ingredient) per acre in a minimum of 5 gallons of water.

For Cooley Spruce Adelgid, apply when white cottony tufts appear.

For Douglas Fir Needle Midge, apply in late April or early May, or before bulbs open.

For Eriophyid needle mite, apply when new growth is 1 inch or less.

As a precaution, treat a few plants of each variety that are going to be sprayed for potential phytotoxicity problems before treating the whole acreage.

**RESTRICTIONS**

- Do not use this product through any type of irrigation system.

24(c) Registrant:  
 Makhteshim-Agan of North America, Inc.  
 551 Fifth Avenue, Suite 1100  
 New York, NY 10176

® denotes registered trademark